

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/44, 31/365, 9/107, 9/48, 47/18		A1	(11) International Publication Number: WO 99/06044 (43) International Publication Date: 11 February 1999 (11.02.99)
(21) International Application Number: PCT/US98/14817 (22) International Filing Date: 27 July 1998 (27.07.98) (30) Priority Data: 60/054,012 29 July 1997 (29.07.97) US (71) Applicant (for all designated States except US): PHARMACIA & UPJOHN COMPANY [US/US]; 301 Henrietta Street, Kalamazoo, MI 49001 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): MOROZOWICH, Walter [US/US]; 5330 Chickadee, Kalamazoo, MI 49002 (US). GAO, Ping [CN/US]; 7191 Crown Pointe Circle, Portage, MI 49024 (US). (74) Agent: YANG, Lucy, X.; Intellectual Property Legal Services, Pharmacia & Upjohn Company, 301 Henrietta Street, Kalamazoo, MI 49001 (US).			(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: PHARMACEUTICAL COMPOSITION FOR ACIDIC LIPOPHILIC COMPOUNDS IN A FORM OF A SELF-EMULSIFYING FORMULATION			
(57) Abstract <p>The present invention provides a novel pharmaceutical composition based on the use of a particular amount of basic amine which comprises a pyranone compound as a pharmaceutically active agent, a basic amine in an amount of from about 0.1 % to about 10 % by weight of the total composition, one or more pharmaceutically acceptable solvents, and one or more pharmaceutically acceptable surfactants. In addition, the composition may further comprise one or more pharmaceutically acceptable oils. The composition is in a form of self-emulsifying formulation which provides high concentration and high oral bioavailability for lipophilic pyranone compounds.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

PHARMACEUTICAL COMPOSITION FOR ACIDIC LIPOPHILIC COMPOUNDS
IN A FORM OF A SELF-EMULSIFYING FORMULATION


FIELD OF THE INVENTION

5 The present invention relates to novel pharmaceutical compositions in a form of a self-emulsifying formulation containing a basic amine which provide high concentration and high oral bioavailability for pyranone compounds which are inhibitors of retroviral protease.

10 BACKGROUND OF THE INVENTION

It has recently been discovered that certain pyranone compounds inhibit retroviral protease and thus they are useful for treating patients infected with human immunodeficiency virus (HIV) which results in acquired immunodeficiency syndrome (AIDS). In particular, the pyranone compound of formula I has been
15 found to be especially effective as an inhibitor of retroviral protease.

20



I

25 However, like many other HIV protease inhibitors, these compounds are characteristically lipophilic and thus poorly water soluble. For example, the compound of formula I has an aqueous solubility about 1 µg/ml in the buffer of pH 6.5 (close to the pH of the intestine), which is considered to have extremely poor aqueous solubility and would be expected to provide very low oral bioavailability in
30 the free acid form. It is well known that an active drug substance or therapeutic moiety administered by any route must possess some aqueous solubility for systemic absorption and therapeutic response. Poorly water soluble compounds often exhibit either incomplete or erratic absorption and thus produce a minimal response at desired dosage.

35 Attempts were made to identify salts of the pyranone compounds in solid forms which could improve aqueous solubility. An overriding defect which has

however remained is that the formulation in the form of salt are prone to precipitation of the parent free acid in the gastrointestinal tract and hence are not capable to provide a dosage in the desired high concentration to permit convenient use and yet meet the required criteria in terms of bioavailability.

5 Recognizing the problems, the present invention is directed toward pharmaceutical compositions in a form of self-emulsifying formulations which provide high concentration and high oral bioavailability for pyranone compounds. In particular it has been discovered that the compositions of the present invention allow the preparation of self-emulsifying formulations containing a pyranone inhibitor of retroviral protease in an exceedingly high concentration up to about 500 mg/g to permit convenient oral administration while at the same time achieving improved bioavailability, which is at least three fold higher than the aqueous suspension of the free acid.

15 INFORMATION DISCLOSURE

The International Publication No. WO 95/30670 discloses pyranone compounds useful to treat retroviral infections.

The International Publication No. WO 96/39142 discloses compositions which increase the bioavailability of protease inhibitors.

20 UK Patent Application, GB 2,222,770A discloses pharmaceutical compositions comprising a cyclosporin in microemulsion pre-concentrate and microemulsion form.

UK Patent Application, GB 2,228,198A discloses pharmaceutical compositions comprising a cyclosporin as active ingredient, a fatty acid triglyceride, a glycerol fatty acid partial ester or propylene glycol or sorbitol complete or partial ester and a tenside having an HLB of at least 10.

UK Patent, GB 2,257,359B discloses pharmaceutical compositions suitable for oral administration comprising a cyclosporin, 1,2-propylene glycol, a mixed mono-, di-, and tri-glyceride and a hydrophilic surfactant.

U.S. Patent No. 4,230,702 discloses a readily enterally absorbable
30 pharmaceutical composition of pharmacologically active agents, which per se are
poorly enterally absorbable.

U.S. Patent Nos. 5,071,643, 5,360,615 and 5,376,688 disclose highly concentrated solutions of an acidic pharmaceutical agent suitable for filling soft gels comprising the acidic pharmaceutical agent and a solvent system, the solvent system comprising polyethylene glycol, water and an ionic species.

SUMMARY OF THE INVENTION

One object of the present invention is to provide a pharmaceutical composition comprising a pyranone compound of formulas I, II, III or IV which provides high oral bioavailability.

- 5 A further object of the present invention is to provide a pharmaceutical composition containing a high drug load of a pyranone compound of formulas I, II, III or IV for convenient administration.

- Another object of the present invention is to provide pharmaceutical compositions which exhibit adequate physical and chemical stability in a self-emulsifying formulation.

Still another object of the present invention is to provide a liquid composition for soft elastic capsules.

- The objects of the present invention have been accomplished in that the present invention provides pharmaceutical compositions in a form of a self-emulsifying formulation which allow a high loading of pyranone compounds (up to about 500 mg/g) while at the same time achieving good oral bioavailability.

The present invention specifically provides a pharmaceutical composition based on the use of a particular amount of a basic amine which comprises:


- (a) a pyranone compound of formula I, II, III or IV,
- 20 (b) a basic amine in an amount of from about 0.1% to about 10% by weight of the total composition,
- (c) one or more pharmaceutically acceptable solvents, and
- (d) one or more pharmaceutically acceptable surfactants.

- In addition, the composition may further comprise one or more pharmaceutically acceptable oils.

DETAILED DESCRIPTION OF THE INVENTION

- In accordance with the present invention, there are pharmaceutical compositions comprising a pyranone compound as a pharmaceutically active agent in a self-emulsifying formulation vehicle.


For the purpose of the present invention, the term "pyranone compounds" refers to compounds of formula II




II

5 wherein R_1 is H-, R_2 is C_3 - C_5 alkyl, phenyl- $(CH_2)_2$ -, het- SO_2NH -(CH_2) $_2$ -, cyclopropyl- $(CH_2)_2$ -, F-phenyl- $(CH_2)_2$ -, het- SO_2NH -phenyl-, or F_3C -(CH_2) $_2$ -; or R_1 and R_2 taken together are a double bond; R_3 is R_4 -(CH_2) $_n$ -CH(R_5)-, H_3C -[O(CH_2) $_2$] $_2$ -CH $_2$ -, C_3 - C_5 alkyl, phenyl- $(CH_2)_2$ -, het- SO_2NH -(CH_2) $_2$ -, (HOCH $_2$) $_3C$ -NH-C(O)-NH-
 10 (CH_2) $_3$ -, (HO $_2C$)(H $_2N$)CH-(CH_2) $_2$ -C(O)-NH-(CH_2) $_3$ -, piperazin-1-yl-C(O)-NH-(CH_2) $_3$ -, HO $_3S$ (CH_2) $_2$ -N(CH_3)-C(O)-(CH $_2$) $_6$ -C(O)-NH-(CH_2) $_3$ -, cyclopropyl- $(CH_2)_2$ -, F-phenyl- $(CH_2)_2$ -, het- SO_2NH -phenyl, or F_3C -(CH_2) $_2$ -; n is 0, 1 or 2; R_4 is phenyl, het, cyclopropyl, H_3C -[O(CH_2) $_2$] $_2$ -, het- SO_2NH -, Br-, N_3 -, or HO $_3S$ (CH_2) $_2$ -N(CH_3)-C(O)-(CH $_2$) $_6$ -C(O)-NH-; R_5 is -CH $_2$ -CH $_3$, or -CH $_2$ -cyclopropyl; R_6 is cyclopropyl,
 15 CH $_3$ -CH $_2$ -, or t-butyl; R_7 is -NR $_8SO_2$ -het, -NR $_8SO_2$ -phenyl, optionally substituted with R_9 , -CH $_2$ -SO $_2$ -phenyl, optionally substituted with R_9 , or -CH $_2$ -SO $_2$ -het; R_8 is -H, or -CH $_3$; R_9 is -CN, -F, -OH, or -NO $_2$; wherein het is a 5-, 6- or 7-membered saturated or unsaturated ring containing from one to three heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur; and including any bicyclic
 20 group in which any of the above heterocyclic rings is fused to a benzene ring or another heterocycle, optionally substituted with -CH $_3$, -CN, -OH, -C(O)OC $_2$ H $_5$, -CF $_3$, -NH $_2$, or -C(O)-NH $_2$; or a pharmaceutically acceptable salt thereof. The preferred compound of formula II is a compound of formula I.

25 The term "pyranone compounds" also refers to compounds of formula III and formula IV



III



IV

30 wherein R_{10} is H-, CH $_3$ O-, or CH $_3$ O-[(CH $_2$) $_2$ O] $_3$ -; R_{11} is cyclopropyl, or -CH $_2$ -CH(CH $_3$) $_2$; R_{12} is -NR $_{14}SO_2$ -phenyl, optionally substituted with R_{15} , -NR $_{14}SO_2$ -het, -CH $_2$ -SO $_2$ -phenyl, optionally substituted with R_{15} , or -CH $_2$ -SO $_2$ -het;
 35 R_{13} is -H, -(CH $_2$) $_2$ -CH $_3$, -CH $_2$ -cyclopropyl, or -CH $_2$ -phenyl; R_{14} is -H, or -CH $_3$; R_{15} is -CN, -F, -CH $_3$, -COOH, or -OH; het is a 5-, 6- or 7-membered saturated or

unsaturated ring containing from one to three heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur; and including any bicyclic group in which any of the above heterocyclic rings is fused to a benzene ring or another heterocycle; optionally substituted with one or two -CH₃, -CN, -C(O)OC₂H₅, or -OH; or a
5 pharmaceutically acceptable salt thereof.

These compounds inhibit retroviral protease and thus inhibit the replication of the virus. They are useful for treating patients infected with human retrovirus such as human immunodeficiency virus (strains of HIV-1 or HIV-2) or human T-cell leukemia viruses (HTLV-I or HTLV-II) which results in acquired immunodeficiency
10 syndrome (AIDS) and/or related diseases. The compounds of formulas I, II, III and IV are disclosed and claimed in International Application No. PCT/US95/05219, incorporated herein by reference, and may be prepared according to the procedures described in International Publication No. WO 95/30670. In particular, the pyranone compound of formula I has been found to be especially effective as an
15 inhibitor of retroviral protease.

The term "self-emulsifying formulation" used herein refers to a concentrated composition capable of generating emulsions or microemulsions upon mixing with sufficient aqueous media.

The emulsions or microemulsions generated from the present invention are
20 conventional solutions comprising a hydrophilic phase and a lipophilic phase. Microemulsions are also characterized by their thermodynamic stability, optical transparency and small average droplet size, generally less than about 0.15 micron.

The term "self-emulsifying formulation vehicle" refers to a composition comprising a basic amine in an amount of from about 0.1% to about 10% by weight
25 of the total composition, one or more pharmaceutically acceptable solvents, and one or more pharmaceutically acceptable surfactants. Optionally, the self-emulsifying formulation vehicle may further comprise one or more pharmaceutically acceptable oils.

The amount of active ingredient in the composition may vary or be adjusted
30 widely depending on the intended route of administration, the potency of the particular active ingredient being used, the severity of the retroviral infection and the required concentration. If desired, however, a pyranone compound as an inhibitor of retroviral protease can be present in the self-emulsifying vehicle of the present invention in an amount up to about 500 mg/g with excellent dispersibility
35 and high oral bioavailability *in vivo* typically reaching 60-96% in rats.

The compositions of the present invention with high oral bioavailability (80-

96% in rats) demonstrate an almost transparent or translucent solution upon dilution with water, which indicates that a microemulsion is formed.

The compositions of the present invention with moderately high bioavailability (60-70% in rats) usually show a visible fine white emulsion without precipitation of the drug upon dilution with water, which indicates that an emulsion is formed.

In one aspect, the present invention specifically provides a pharmaceutical composition based on the use of particular oil phase which comprises:

- (a) a pyranone compound of formulas I, II, III or IV as a pharmaceutically active agent,
- (b) a basic amine in an amount of from about 0.1% to about 10% by weight of the total composition,
- (c) one or more pharmaceutically acceptable solvents, and
- (d) one or more pharmaceutically acceptable surfactants.

In addition, the composition may further comprise one or more pharmaceutically acceptable oils.

The term "pharmaceutically acceptable" used herein refers to those properties which are biologically compatible with the treated subjects from a pharmacological and toxicological point of view.

Solvents of the present invention refer to propylene glycol, polypropylene glycol, polyethylene glycol (such as PEG300, 400, 600, etc.), glycerol, ethanol, triacetin, dimethyl isosorbide, glycofurol, propylene carbonate, water, dimethyl acetamide or a mixture thereof.

The preferred solvent is propylene glycol or a mixture comprising propylene glycol and 95% (v/v) ethanol (hereinafter ethanol). In the mixture of propylene glycol and ethanol, propylene glycol is in an amount of from about 50% to about 95%.

Surfactants of the present invention refer to non-ionic surfactants including Polyoxyl 40 hydrogenated castor oil sold under the trade name, among the others, Cremophor RH40; Polyoxyl 35 castor oil sold under the trade name, among the others, Cremophor EL or Cremophor EL; Polysorbates; Solutol HS-15; Tagat TO; Peglicol 6-oleate; Polyoxyethylene stearates; Saturated Polyglycolized Glycerides; or Poloxamers; all of which are commercially available. The preferred surfactant is Cremophor RH40, Cremophor EL or Polysorbate 80.

Saturated Polyglycolized Glycerides used herein include Gelucire 44/14 or Gelucire 50/13.

Polyoxyethylene stearates used herein include Poloxyl 6 stearate, Poloxyl 8 stearate, Poloxyl 12 stearate and Poloxyl 20 stearate.

Poloxamers used herein include Poloxamer 124, Poloxamer 188, Poloxamer 237, Poloxamer 338 and Poloxamer 407.

5 Polysorbates used herein include Polysorbate 20, Polysorbate 40, Polysorbate 60 and Polysorbate 80.

The term "basic amine" used herein refers to lower alkylamine such as, for example, ethanolamine, diethanolamine, triethanolamine, dimethylaminoethanol, tris(hydroxymethyl)aminomethane or ethylenediamine; quaternary ammoniums such
10 as, for example, choline hydroxide; basic amino acids such as, for example, arginine lysine or guanidine. The preferred basic amine is lower alkylamine or quaternary ammonium. The preferred lower alkylamine is dimethylaminoethanol or tris(hydroxymethyl)aminomethane.

Oil useful in forming the composition of the present invention includes a
15 broad spectrum of water-immiscible materials such as soybean oil, avocado oil, squalene oil, sesame oil, olive oil, canola oil, corn oil, rapeseed oil, safflower oil, sunflower oil, fish oils, flavored oils, water insoluble vitamins and mixtures thereof. The preferred oils are the medium chain fatty acid glyceride such as commercially available under the trade names, among the others, Maisine, Miglyol 812, Captex
20 355, Myritol, Capmul MCM, Captex 200, Myvacet, Myverol 18-92, Arlacel 186, Neobee, Mazol or commercially available monoolein, diolein and triolein. The more preferred oils of the present invention is Capmul MCM, monoolein, diolein, triolein, monolinoleate, dilinoleate, trilinoleate or Maisine.

A typical composition according to the present invention comprises:

- 25 (a) a pyranone compound of formulas I, II, III or IV in an amount of from about 1% to about 50% by weight of the total composition,
(b) a basic amine in an amount of from about 0.1% to about 10% by weight of the total composition,
(c) one or more pharmaceutically acceptable solvents in an amount of from about
30 10% to about 30% by weight of the total composition, and
(d) a pharmaceutically acceptable surfactant in an amount of from about 10% to about 50% by weight of the total composition.

Optionally, the composition further comprises one or more oils in an amount of from about 5% to 35% by weight of the total composition.

35 A preferred composition of the present invention comprises:

- (a) a pyranone compound of formulas I, II, III or IV in an amount of from about

20% to about 30% by weight of the total composition,

- (b) a dimethylaminoethanol or tris(hydroxymethyl)aminomethane in an amount of from about 0.1% to about 7% by weight of the total composition,
- (c) a solvent comprising propylene glycol or a mixture of propylene glycol and ethanol in an amount of from about 15% to about 25% by weight of the total composition, and
- (d) a surfactant comprising Polysorbate 80 in an amount of from about 30% to about 45% by weight of the total composition .

Another preferred composition of the present invention comprises:

- (a) a pyranone compound of formulas I, II, III or IV in an amount of from about 20% to about 30% by weight of the total composition,
- (b) a dimethylaminoethanol or tris(hydroxymethyl)aminomethane in an amount of from about 0.1% to about 7% by weight of the total composition,
- (c) a solvent comprising propylene glycol or a mixture of propylene glycol and ethanol in an amount of from about 15% to about 25% by weight of the total composition,
- (d) a surfactant comprising Cremophor RH40 or Cremophor EL in an amount of from about 30% to about 45% by weight of the total composition, and
- (e) one or more oils selected from the group consisting of monoolein, diolein, Campul MCM or Maisine in an amount of from about 5% to about 25% by weight of the total composition.

Optionally, the preferred compositions further comprise one or more oils in an amount of about 5% to about 25% by weight of the total composition.

In the preferred compositions of the present invention, the mixture of propylene glycol and ethanol is in a ratio of about 1:1.

In the preferred compositions of the present invention, an even more preferred composition comprises an dimethylaminoethanol or tris(hydroxymethyl)aminomethane in an amount of from about 0.1% to 5% by weight of the total composition.

In particular, the most preferred composition of the present invention comprises the pyranone compound of formula I.

The composition of the present invention may take the form of liquid for soft elastic capsules or hard gelatin capsules by oral application. The composition may also be in the form of liquid solution for oral, parenteral, rectal or topical application. The preferred dosage form is in the form of liquid for soft elastic capsules.

If desired, the compositions of the present invention may further comprise conventional pharmaceutical additives such as co-surfactant (for example, sodium lauryl sulfate), coloring agents, flavoring agents, fragrances, preserving agents, stabilizers, anti-oxidant and/or thickening agents.

5 The compositions of the present invention may be prepared in a conventional manner, for example, by dissolving an active agent in the solvent, then adding the basic amine, the surfactant and optionally an oil. The resulting solution is then formulated into the desired dosage form such as, for example, soft elastic capsules or hard gelatin capsules by known manufacturing technology.

10 The pharmaceutical compositions of the present invention will be better understood in connection with the following examples, which are intended as an illustration of and not a limitation upon the scope of the invention. Without further elaboration, it is believed that one skilled in the art can, using the preceding description and the information provided in the examples below, practice the present
15 invention to its fullest extent.

A. General Procedure for Preparing the Compositions of the Present Invention.

Drug is placed in a container. A solvent comprising propylene glycol or a mixture of solvents selected from ethanol (95%) and propylene glycol (1:1 by weight) is added and the cap is tightened. The container is put in a water bath at about 60
20 °C and shaken gently until all of the drug material is dissolved. After the container is cooled down to room temperature, appropriate amounts of a basic amine, a surfactant and optionally one or more oils are added into the container. The container is sealed and put in a water bath at about 60 °C and shaken gently until a clear solution is formed. The container is usually left at ambient conditions for
25 future use.

EXAMPLE 1

Component	Weight (mg)	%, w/w
The compound of formula I	299	29.8
EtOH/Propylene Glycol (1:1)	250	25.0
30 Polysorbate 80	375	37.5
Ethanolamine	60	6.0
Sodium lauryl sulfate	20	2.0

EXAMPLE 2

(The same formulation as the EXAMPLE 1 except without ethanolamine)

5

Component	Weight (mg)	%, w/w
The compound of formula I	302	31.3
EtOH/Propylene Glycol (1:1)	258	26.8
Polysorbate 80	385	39.9
Sodium lauryl sulfate	19	2.00

EXAMPLE 3

10

Component	Weight (mg)	%, w/w
The compound of formula I	298	28.0
EtOH/Propylene Glycol (1:1)	200	18.6
Cremophor EL	505	47.5
Dimethylaminoethanol	63	5.8

15

EXAMPLE 4

20

Component	Weight (mg)	%, w/w
The compound of formula I	296	29.6
EtOH/Propylene Glycol (1:1)	258	25.8
Polysorbate 80	358	35.8
Dimethylaminoethanol	70	7.0
Sodium lauryl sulfate	18	1.8

25

EXAMPLE 5

5	Component	Weight (mg)	%, w/w
	The compound of formula I	303	29.5
	EtOH/Propylene Glycol (1:1)	200	19.4
	Campul MCM	65	6.3
	Cremophor EL	450	43.3
	Diethanolamine	15	1.4

EXAMPLE 6

10	Component	Weight (mg)	%, w/w
	The compound of formula I	301	26.2
	EtOH/Propylene Glycol (1:1)	203	17.7
	Monoolein	252	22.0
	Cremophor RH40	312	27.2
	Ethanolamine	62	5.4
15	Sodium lauryl sulfate	16	1.4

EXAMPLE 7

20	Component	Weight (mg)	%, w/w
	The compound of formula I	302	26.2
	EtOH/Propylene Glycol (1:1)	203	17.6
	Diolein	264	22.9
	Cremophor RH40	305	26.4
	Ethanolamine	64	5.6
25	Sodium lauryl sulfate	15	1.3

EXAMPLE 8

	Component	Weight (mg)	%, w/w
	The compound of formula I	303	29.5
	EtOH/Propylene Glycol (1:1)	200	19.4
5	Campul MCM	65	6.3
	Cremophor EL	450	43.3
	Dimethylaminoethanol	15	1.4

EXAMPLE 9

10	Component	Weight (mg)	%, w/w
	The compound of formula I	303	29.5
	EtOH/Propylene Glycol (1:1)	200	19.4
	Campul MCM	65	6.3
	Cremophor EL	450	43.3
15	Choline hydroxide	15	1.4

EXAMPLE 10

	Component	Weight (mg)	%, w/w
	The compound of formula I	400	39.90
20	EtOH (95% pure)	100	9.97
	Propylene Glycol	40	3.99
	Propyl gallate	2.5	2.49
	Cremophor EL	360	35.91
	Diethanolamine	80	7.98
25	H ₂ O	20	1.99

EXAMPLE 11

Component	Weight (mg)	%, w/w
The compound of formula I	500	49.87
EtOH (95%)	90	8.98
Propyl gallate	2.5	0.25
Cremophor EL	320	31.92
Ethanolamine	90	8.98

B. Oral Bioavailability Test.

10 (i) Oral Bioavailability Test in Rats

Sprague-Dawley male rats were selected for the *in vivo* oral bioavailability study. Each rat was prepared by the surgical implantation of an indwelling cannula in the superior vena cava. Each rat, in the weight range of 300 - 400 g, was fasted overnight prior to dosing. Each formulation was orally administered to a group of rats (n=3) at a 20 mg/kg dose. The formulations with high concentration of the compound of formula I (typically 200-300 mg/g) was diluted by 100-fold with water and injected directly into the rat's stomach using oral gavage. Serial blood samples of 0.25 ml were obtained from the indwelling cannula at 0.25, 0.5, 1, 2, 4, 6, 8, 12, and 24 hours after dosing. These blood samples were analyzed using a HPLC assay specific for the compound of formula I. Drug concentrations in the blood of the test rats are plotted against the time after the drug is administered through an intravenous (i.v.) or oral route and the AUCs (the Area Under the plasma concentration-time Curve) are integrated using the trapezoidal rule to calculate the absolute bioavailability.

$$\text{Absolute bioavailability (F)} = \frac{(AUC)_{oral} / Dose_{oral}}{(AUC)_{iv} / Dose_{iv}}$$

The present invention achieves the desired results as demonstrated in the absolute bioavailability test in Table 1. Comparing the reference example 2 (without basic amine) with the examples 1, 5, 6 and 7, the formulations of the present invention exhibit at least three times fold higher oral bioavailability.

TABLE 1
Oral Bioavailability Test in Rats

	Example No.	Absolute Mean Oral Bioavailability (%)
	1	65
5	2	17
	5	96
	6	66
	7	58
10	Aqueous suspension of free acid of the compound of formula I	< 20


15

We claim

1. A pharmaceutical composition comprising:

(a) a pyranone compound of formula II as a pharmaceutically active agent,

5



II

10 (b) a basic amine in an amount of from about 0.1% to about 10% by weight of the total composition,

(c) one or more pharmaceutically acceptable solvents, and

(d) one or more pharmaceutically acceptable surfactants;

wherein R_1 is H-;

15 R_2 is C_3 - C_5 alkyl, phenyl- $(CH_2)_2$ -, het-SO₂NH- $(CH_2)_2$ -, cyclopropyl- $(CH_2)_2$ -, F-phenyl- $(CH_2)_2$ -, het-SO₂NH-phenyl-, or F_3C -(CH_2)₂-; or

R_1 and R_2 taken together are a double bond;

R_3 is R_4 -(CH_2)_n-CH(R_5)-, H_3C -[O(CH_2)₂]₂-CH₂-, C_3 - C_5 alkyl, phenyl- $(CH_2)_2$ -, het-SO₂NH- $(CH_2)_2$ -, (HOCH₂)₃C-NH-C(O)-NH- $(CH_2)_3$ -,

20 (HO₂C)(H₂N)CH- $(CH_2)_2$ -C(O)-NH- $(CH_2)_3$ -, piperazin-1-yl-C(O)-NH- $(CH_2)_3$ -, HO₃S(CH₂)₂-N(CH₃)-C(O)- $(CH_2)_6$ -C(O)-NH- $(CH_2)_3$ -, cyclopropyl- $(CH_2)_2$ -, F-phenyl- $(CH_2)_2$ -, het-SO₂NH-phenyl, or F_3C -(CH_2)₂-;

n is 0, 1 or 2;

R_4 is phenyl, het, cyclopropyl, H_3C -[O(CH_2)₂]₂-, het-SO₂NH-, Br-, N₃-, or

25 HO₃S(CH₂)₂-N(CH₃)-C(O)- $(CH_2)_6$ -C(O)-NH-;

R_5 is -CH₂-CH₃, or -CH₂-cyclopropyl;

R_6 is cyclopropyl, CH₃-CH₂-, or t-butyl;

R_7 is -NR₈SO₂-het, -NR₈SO₂-phenyl, optionally substituted with R_9 , -CH₂-SO₂-phenyl, optionally substituted with R_9 , or -CH₂-SO₂-het;

30 R_8 is -H, or -CH₃;

R_9 is -CN, -F, -OH, or -NO₂;

wherein het is a 5-, 6- or 7-membered saturated or unsaturated ring containing from one to three heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur; and including any bicyclic group in which any of the above heterocyclic rings


35 is fused to a benzene ring or another heterocycle, optionally substituted with -CH₃, -CN, -OH, -C(O)OC₂H₅, -CF₃, -NH₂, or -C(O)-NH₂; or

a pharmaceutically acceptable salt thereof.

2. The pharmaceutical composition of claim 1 wherein the pyranone compound of formula II is a compound of formula I.

5

10




I.


15 3. A pharmaceutical composition comprising:

(a) a pyranone compound of formula III or IV as a pharmaceutically active agent,

20



III



IV

25 (b) a basic amine in an amount of from about 0.1% to about 10% by weight of the total composition,

(c) one or more pharmaceutically acceptable solvents, and

(d) one or more pharmaceutically acceptable surfactants;

wherein R_{10} is H-, CH_3O -, or $\text{CH}_3\text{O}-[(\text{CH}_2)_2\text{O}]_3$ -;

R_{11} is cyclopropyl, or $-\text{CH}_2-\text{CH}(\text{CH}_3)_2$;

30 R_{12} is $-\text{NR}_{14}\text{SO}_2$ -phenyl, optionally substituted with R_{15} , $-\text{NR}_{14}\text{SO}_2$ -het, $-\text{CH}_2-\text{SO}_2$ -phenyl, optionally substituted with R_{15} , or $-\text{CH}_2-\text{SO}_2$ -het;

R_{13} is H-, $-(\text{CH}_2)_2-\text{CH}_3$, $-\text{CH}_2$ -cyclopropyl, or $-\text{CH}_2$ -phenyl;

R_{14} is H-, or $-\text{CH}_3$; R_{15} is $-\text{CN}$, $-\text{F}$, $-\text{CH}_3$, $-\text{COOH}$, or $-\text{OH}$;

35 het is a 5-, 6- or 7-membered saturated or unsaturated ring containing from one to three heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur, and including any bicyclic group in which any of the above heterocyclic rings is fused

to a benzene ring or another heterocycle; optionally substituted with one or two -CH₃, -CN, -C(O)OC₂H₅, or -OH; or a pharmaceutically acceptable salt thereof.

4. The pharmaceutical composition of claim 1 or 3 wherein a compound of
5 formula II, III or IV is in an amount of from about 1% to about 50%.
5. The pharmaceutical composition of claim 2 wherein a compound of formula I is in an amount of from about 20% to about 30%.
- 10 6. The pharmaceutical composition of claim 1 or 3 wherein the basic amine is lower alkylamine, basic amino acid, or choline hydroxide.
7. The pharmaceutical composition of claim 6 wherein the lower alkylamine is ethanolamine, diethanolamine, triethanolamine, tris(hydroxymethyl)aminomethane,
15 ethylenediamine, or dimethylaminoethanol.
8. The pharmaceutical composition of claim 6 wherein the lower alkylamine is dimethylaminoethanol or tris(hydroxymethyl)aminomethane.
- 20 9. The pharmaceutical composition of claim 6 wherein the basic amino acid is arginine, lysine or guanidine.
10. The pharmaceutical composition of claim 1 or 3 wherein the basic amine is in an amount of from about 0.1% to about 7% by weight of the total composition.
25
11. The pharmaceutical composition of claim 1 or 3 wherein the basic amine is in an amount of from about 0.1% to about 5% by weight of the total composition.
12. The pharmaceutical composition of claim 1 or 3 wherein the pharmaceutically
30 acceptable solvent is propylene glycol, polypropylene glycol, polyethylene glycol, glycerol, ethanol, triacetin, dimethyl isosorbide, glycofurool, propylene carbonate, water, dimethyl acetamide or a mixture thereof.
13. The pharmaceutical composition of claim 1 or 3 wherein the pharmaceutically
35 acceptable solvent is propylene glycol.

14. The pharmaceutical composition of claim 1 or 3 wherein the pharmaceutically acceptable solvent is a mixture solution comprising propylene glycol and 95% (v/v) ethanol in a ratio of about 1:1.
- 5 15. The pharmaceutical composition of claim 1 or 3 wherein the pharmaceutically acceptable solvent is in an amount of from about 10% to about 30% by weight of the total composition.
16. The pharmaceutical composition of claim 1 or 3 wherein the pharmaceutically acceptable solvent is in an amount of from about 15% to about 25% by weight of the total composition.
- 10 17. The pharmaceutical composition of claim 1 or 3 wherein the pharmaceutically acceptable surfactant is Polyoxyl 40 hydrogenated castor oil, Polyoxyl 35 castor oil, Polysorbate 20, Polysorbate 40, Polysorbate 60, Polysorbate 80, Solutol HS-15, Tagat TO, Peglicol 6-oleate, Polyoxyethylene stearates, Saturated Polyglycolized Glycerides, or Poloxamers.
18. The pharmaceutical composition of claim 1 or 3 wherein the pharmaceutically acceptable surfactant is Polyoxyl 40 hydrogenated castor oil, Polyoxyl 35 castor oil, or Polysorbate 80.
- 20 19. The Polyoxyl 40 hydrogenated castor oil of claim 18 which is Cremophor RH40.
- 25 20. The Polyoxyl 35 hydrogenated castor oil of claim 18 which is Cremophor EL or Cremophor EL-P.
21. The pharmaceutical composition of claim 1 or 3 wherein the surfactant is in an amount of from about 10% to about 50% by weight of the total composition.
- 30 22. The pharmaceutical composition of claim 1 or 3 wherein the surfactant is in an amount of from about 30% to about 45% by weight of the total composition.
- 35 23. The pharmaceutical composition of claim 1 or 3 wherein the composition further comprises one or more oils.

24. The pharmaceutical composition of claim 23 wherein said oil is soyb an oil, avocado oil, squalene oil, sesame oil, olive oil, canola oil, corn oil, rapeseed oil, safflower oil, sunflower oil, fish oils, flavored oils, water insoluble vitamins or a mixture thereof.
- 5
25. The pharmaceutical composition of claim 23 wherein said oil is monoolein, diolein, triolein, monolinoleate, dilinoleate, trilinoleate, Maisine, Miglyol 812, Captex 355, Myritol, Capmul MCM, Captex 200, Myvacet, Myverol 18-92, Arlacel 186, Neobee or Mazol.
- 10
26. The pharmaceutical composition of claim 25 wherein said oil is Capmul MCM, monoolein, diolein, monolinoleate, dilinoleate or Maisine.
27. The pharmaceutical composition of claim 23 wherein said oil is in an amount
15 from about 5% to about 35% by weight of the total composition.
28. A pharmaceutical composition comprising:
- (a) a pyranone compound of formulas I in an amount of from about 20% to about 30% by weight of the total composition,
- 20 (b) a dimethylaminoethanol or tris(hydroxymethyl)aminomethane in an amount of from about 0.1% to about 5% by weight of the total composition,
- (c) a solvent comprising propylene glycol or a mixture of propylene glycol and 95% (v/v) ethanol in an amount of from about 15% to about 25% by weight of the total composition, and
- 25 (d) a surfactant comprising Polysorbate 80 in an amount of from about 30% to about 45% by weight of the total composition.
29. A pharmaceutical composition comprising:
- (a) a pyranone compound of formulas I in an amount of from about 20% to about
30 30% by weight of the total composition,
- (b) a dimethylaminoethanol or tris(hydroxymethyl)aminomethane in an amount of from about 0.1% to about 5% by weight of the total composition,
- (c) a solvent comprising propylene glycol or a mixture solution of propylene glycol and 95% (v/v) ethanol in an amount of from about 15% to about 25%
35 by weight of the total composition,
- (d) a surfactant comprising Cremophor RH40 or Cremophor EL in an amount of

- from about 30% to about 45% by weight of the total composition, and
- (e) an oil comprising monoolein, diolein, monolinoleate, dilinoleate Campul MCM or Maisine in an amount of from about 5% to about 25% by weight of the total composition.

5

30. The pharmaceutical composition of claim 28 or 29 wherein the mixture solution of propylene glycol and ethanol is in a ratio of about 1:1.

31. The pharmaceutical composition of claim 1, 3, 28 or 29 which is a self-emulsifying formulation capable of generating emulsions or microemulsions upon mixing with sufficient aqueous media.

32. The pharmaceutical composition of claim 1, 3, 28 or 29 which is in a form of liquid for soft elastic capsules.

15

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 98/14817

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K31/44 A61K31/365 A61K9/107 A61K9/48 A61K47/18

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, A	WO 98 22106 A (ABBOT LABORATORIES) 28 May 1998 see the whole document -----	1-32

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

4 November 1998

Date of mailing of the international search report

12/11/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Ventura Amat, A

Information on patent family members

PCT/US 98/14817

Form PCT/ISA/210 (patent family annex) (July 1992)